

Structure of the GAF Domain, a Ubiquitous Signaling Motif and a new Class of Cyclic GMP Receptor

Y.-S.J. Ho, L. Burden, and J. Hurley (*Laboratory of Molecular Biology, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health*)

Abstract No. Ho9322

Beamline(s): **X9B**

GAF domains are ubiquitous motifs present in cyclic **GMP**-regulated cyclic nucleotide phosphodiesterases, certain **adenylyl** cyclases, the bacterial transcription factor **FhlA**, and hundreds of other signaling and sensory proteins from all three kingdoms of life. The crystal structure of the *Saccharomyces cerevisiae* YKG9 protein was determined at 1.9 Å resolution. The structure revealed a fold that resembles the PAS domain, another ubiquitous signaling and sensory transducer. YKG9 does not bind cyclic GMP, but the isolated first GAF domain of phosphodiesterase 5 binds with $K_d = 650$ nM. The cyclic GMP binding site of the phosphodiesterase GAF domain was identified by homology modeling and site-directed mutagenesis, and consists of conserved Arg, Asn, Lys, and Asp residues. The structural and binding studies taken together show that the cyclic GMP-binding GAF domains form a new class of cyclic nucleotide receptors distinct from the regulatory domains of cyclic nucleotide-regulated protein kinases and ion channels.